

# Comparative study on morbidity pattern among adolescent schoolgirls in rural and urban school of Jhansi

Vimal Arya, Shobha Chaturvedi

Department of Community Medicine, Maharani Laxmi Bai Medical College, Jhansi, Uttar Pradesh, India

Correspondence to: Vimal Arya, E-mail: drvimalarya@gmail.com

Received: January 16, 2018; Accepted: February 12, 2018

## ABSTRACT


**Background:** The World Health Organization (WHO) proposed that the age limit of 10–19 years be used to identify as the adolescent age. According to the WHO, adolescents constitute about one-fifth of the world population, and in India, they constitute about 21% of the total population. Most of the surveys show that health status of adolescent girls is at suboptimal level. **Objectives:** The objectives of this study were to assess nutritional status and morbidity pattern among the adolescent girls and to suggest measures for the improvement of health status of adolescent girls. **Materials and Methods:** A community-based cross-sectional study was conducted among 250 adolescent schoolgirls in rural and urban field practice area of Jhansi school from January 2017 to July 2014. **Results:** Among the various morbidities, eye problem was seen in maximum number of adolescent girls. Eye problem was present in 44.8% of adolescent girls followed by respiratory 14.7% and ear 13.06% disease. Skin disease was present in 3.2% of adolescent girls, which was more in rural girls 6.7% than in urban girls 1.7%, may be due to better hygienic practice in urban schoolgirls. **Conclusion:** Rural background, low socioeconomic status, illiteracy, birth rate and order, income, and number of members in a family have shown to be significant determinants of morbidity pattern in the adolescent girls.

**KEY WORDS:** Adolescent; Morbidity Rural; Urban

## INTRODUCTION

The World Health Organization defines adolescence as the segment of life between the ages of 10 and 19 years.<sup>[1,2]</sup> 25% of adult height and up to 50% of adult weight are attained during adolescence.<sup>[3]</sup> It is also an intense anabolic period when requirements for all nutrients increase.<sup>[4]</sup> 85% of them live in developing countries.<sup>[5]</sup> Adolescent girls form an important vulnerable sector of population. Adolescent period is a growth stage of a girl's life and is a unique intervention point in the

lifecycle for a number of reasons; it is a transition phase through which a child becomes an adult, which is accompanied with hormonal and physical changes marked by spurt in growth. In India, adolescents account for 21% (approximately 230 million) of the total population among which the adolescent girls comprise about 10.3% of the total population of India.<sup>[3,4]</sup> The importance of this target group lies in the fact that they are going to be the mothers of tomorrow - whose well-being is critically important for improving the nutritional, health, and educational status of women in the state. Thus, it is a crucial period of women's life where sociocultural factors not only influence her health but also determine the health of future generations. Majority of adolescents still do not have access to information on reproductive health and rights, nor do they have access to preventive and curative services.<sup>[6]</sup> The health problems of adolescents are very different from those of younger children and adults.<sup>[7]</sup> Nutritional deficiency

Access this article online	
Website: <a href="http://www.ijmsph.com">http://www.ijmsph.com</a>	Quick Response code
DOI: 10.5455/ijmsph.2018.0101812022018	

International Journal of Medical Science and Public Health Online 2018. © 2018 Vimal Arya and Shobha Chaturvedi. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material for any purpose, even commercially, provided the original work is properly cited and states its license.

disorders (stunting, wasting), menstrual disorders, and reproductive tract infections/sexually transmitted diseases/HIV/AIDS have been appeared as serious problem during this stage.<sup>[8]</sup> The vicious cycle of undernutrition and its impact on health status of adolescent girls is increasingly being studied in terms of nutritional deficiencies. One of the major impacts of undernutrition and compromised health status of adolescent girls is reflected by high prevalence of anemia. Based on the above issues, the rationale behind this study was to evaluate the status of school going adolescent girls in terms of nutrition and disease who are the most hit age group in the state of Uttar Pradesh.

### Aims and Objectives

To assess nutritional status and morbidity pattern among the adolescent girls and to suggest measures for the improvement of health status of adolescent girls.

### MATERIALS AND METHODS

The present school-based cross-sectional study was conducted for 6 months in Jhansi, from January 2017 to July 2017. Using multistage random sampling technique, four schools were randomly selected in urban area and 176 adolescent girls were chosen from these schools. Similarly, in rural area, two schools were randomly selected and from there 74 adolescent girls were chosen. (In Jhansi District, 70% of the population lives in urban area and 30% of the population lives in rural area).

All adolescent girls registered in school were decided to include in the study after taking verbal consent. Data were collected in pre-designed pro forma by interview technique and clinical examination of the girls in the presence of their mothers. The sociodemographic data of all the adolescent girls were recorded. Biosocial characteristic of adolescent girl was age, birth order, total family member, type of family, religion, caste, education status, education, and occupation of parent's total family income, with the help of these per capita incomes was calculated. The pre-tested questionnaire included clinical examination; history of chronic illness in last 6 months and the symptoms of illness were recorded. Health status of the adolescent girls was assessed by the morbidity pattern.

Data were collected, compiled, and tabulated using Microsoft Excel and analyzed using SPSS 17.0 Version for calculation of percentages.

### RESULTS

A total of 250 school going adolescent girls (176 urban and 74 rural) aged 10–19 years were interviewed. Our study

revealed that the most of rural schoolgirls were of 14 years Table 1.

According to Table 2, majority of adolescent girls were Hindus (72.8%), while Muslim girls were only (27.2%). About 42.8% of the adolescent girls in our study belonged to other backward classes (OBC).

Table 3 summarizes the system wise morbidity pattern in adolescent schoolgirls. 44.8% girls of urban and 22.9% girls of rural schoolgirls were having eye problem. 14.7% girls in urban and 17.5% girls in rural school were having respiratory tract infection. About 11.3% of urban and 16.21% of rural adolescent girls were suffering from reproductive system morbidity.

Table 4 summarizes that there was no association seen between reproductive tract illnesses and place of schools. A total of 10.4% girls had vaginal discharge in which urban was 7.9% and rural was 16.21%.

### DISCUSSION

In the present study, majority (58.4%) of girls were in the age group of 14–16 years and mean age was 14.12. About 72.8% adolescent girls were Hindus. The highest percentage of girls belonged to OBC. Among the various morbidities, eye problem was seen in maximum number of adolescent girls. Eye problem was present in 85.8% of adolescent girls followed by ear (27.8%) and respiratory (15.6%) disease. Skin disease was present in 3.2% of adolescent girls, which was more in rural girls than in urban girls, may be due to better hygienic practice in urban schoolgirls.

However, Baliga *et al.* and Guduri *et al.* in their study reported that the mean age of rural adolescent girls was 12.9 years and 12.67 which was less than our study.<sup>[10,11]</sup> As per National Family Health Survey-3, 82.6% of households in Uttar Pradesh were Hindus and 16.3% Muslims and

**Table 1:** Distribution of adolescent schoolgirls by age

Age (years)	Urban n=176 (%)	Rural n=74 (%)	Total n=250 (%)
10	1 (0.5)	3 (4.05)	4 (1.6)
11	15 (8.5)	5 (6.7)	20 (8)
12	18 (10.2)	5 (6.7)	23 (9.2)
13	10 (5.6)	4 (5.4)	14 (5.6)
14	36 (20.4)	16 (21.6)	52 (20.8)
15	31 (17.6)	16 (21.6)	47 (18.8)
16	34 (19.3)	13 (17.5)	47 (18.8)
17	22 (12.5)	4 (5.4)	26 (10.4)
18	7 (3.9)	5 (6.7)	12 (4.8)
19	2 (1.1)	3 (4.05)	5 (2)

**Table 2:** Distribution of adolescent schoolgirls according to their biosocial characteristics

Biosocial characteristics	Urban n=176 (%)	Rural n=74 (%)	Total n=250
Religion			
Hindu	121 (68.7)	61 (82.4)	182 (72.8)
Muslim	55 (31.2)	13 (17.5)	68 (27.2)
Caste			
General	63 (35.7)	23 (31.08)	86 (34.4)
OBC	71 (40.3)	36 (48.6)	107 (42.8)
SC/ST	42 (23.8)	15 (20.2)	57 (22.8)
Type of family			
Nuclear	166 (94.3)	60 (81.08)	226 (90.4)
Joint	10 (56)	14 (18.9)	24 (9.6)
Total family members			
Up to 5	79 (44.8)	30 (40.5)	109 (43.6)
>5	97 (55.1)	44 (59.4)	141 (56.4)
Birth order			
1-3	117 (66.4)	51 (68.9)	168 (67.2)
3-4	55 (31.25)	19 (25.67)	74 (29.6)
≥5	4 (2.21)	4 (5.4)	8 (3.2)
Socioeconomic status			
I	2 (1.13)	0 (0)	2 (0.8)
II	72 (40.9)	1 (1.35)	73 (29.2)
III	60 (34.09)	43 (58.10)	103 (41.2)
IV	42 (23.8)	30 (40.5)	72 (28.8)
V	-	-	-

OBC: Other backward classes

**Table 3:** Distribution of system wise morbidity pattern in adolescent schoolgirls

Systemic morbidity	Urban n=176 (%)	Rural n=74 (%)	Total n=250 (%)
Skin	3 (1.7)	5 (6.7)	8 (3.2)
Eye	79 (44.8)	17 (22.9)	96 (38.5)
Ear	23 (13.06)	12 (16.21)	35 (14.0)
Respiratory	26 (14.7)	13 (17.5)	39 (15.6)
GIT	17 (3.6)	7 (9.4)	24 (9.6)
Cardiovascular system	8 (4.5)	8 (10.8)	16 (6.4)
Reproductive system	20 (11.3)	12 (16.21)	32 (12.8)

GIT: Gastrointestinal tract

overall 41.9% adolescent girls belonged to OBC. Almost similar finding was seen in a study conducted by Yerpude *et al.* who reported the prevalence of ophthalmic problems (12.5%) and respiratory infection (8.33%).<sup>[12]</sup> Sachan *et al.* reported that skin disease in 3.2% of girls in both the urban and rural schools.<sup>[13]</sup> As compared to system wise morbidity of adolescent girls, Yerpude *et al.* showed highest morbidity

**Table 4:** Distribution of adolescent schoolgirls according to reproductive tract morbidity

Reproductive tract illness	Urban n=176 (%)	Rural n=74 (%)	Total n=250 (%)
Lower abdominal pain	2 (1.13)	0 (0)	2 (1.13)
Vaginal discharge	14 (7.9)	12 (16.21)	26 (10.4)
Others	4 (2.2)	0 (0)	4 (2.2)

of respiratory tract infection (23.5%) as compared to our study which has eye morbidity as common on both rural and urban (44.8%, 22.9%), but the reproductive tract illness was found to be like our study.<sup>[12]</sup> In a study conducted by Sachan *et al.*,<sup>[9]</sup> systemic morbidity was found to be higher in rural area as compared to our study but slightly higher in urban area which depicts the seasonal variation in states of the country.<sup>[13]</sup> According to Guduri *et al.*, skin and eye diseases are common in both rural and urban areas of Chennai (21%, 19%, 17%, and 25%) which signifies the level of hygiene maintained by adolescent girls as compared to the current study.<sup>[11]</sup>

### Strengths and Limitations

Strength of the present study is that it has shown an important public health aspect which are adolescent girls and their morbidity pattern and in the state of Uttar Pradesh which is under Empowered Action Group of India. Second, comparison between urban and rural girls toward morbidity will help in implementing policies and draft for their upliftment in nutritional status. This study has certain limitations; first, number of students in urban and rural are different which can affect the results of the study though it is done by random sampling. Second, pilot testing was not done. Third, investigation reports were not seen for confirmation only verbal reports were recorded.

### CONCLUSION

High morbidity was found in adolescent schoolgirls. Health education regarding reproductive health and its morbidities should be conducted in schools and colleges and in communities. It can be included as a part of school health program. Emphasis should be given to make them aware about the importance of seeking of health care. There is a need for regular supply of iron and folic acid tablets at Anganwadi workers and to increase the compliance regarding consuming tablets among adolescent girls. The health and nutritional status of adolescent girls is very poor in rural areas. It is because of gender discrimination in the families. Regular health check-ups and periodical examination in the schools and families should be done by health workers. Deworming should be done on a mass level at regular intervals. Adolescent clinics should be opened in the health

centers. Further, research can be encouraged to improve the health status of the inmates of the social welfare hostels.

## REFERENCES

1. Adesina AF, Peterside O, Anochie I, Akani NA. Weight status of adolescent in secondary school in port Harcourt using body mass index. *Ital J Pediatr* 2012;38:31.
2. Jain K, Garg SK, Singh JV, Bhatnagar M, Chopra H, Bajpai SK. Reproductive health of adolescent girls in an urban population of Meerut, Uttar Pradesh. *Health Popul Prospect Issues* 2009;32:204-9.
3. WHO. Adolescent Nutrition: A Review of the Situation in Selected South-East Asian Countries. New Delhi: WHO; 2006.
4. DiMeglio G. Nutrition in adolescence. *Pediatr Rev* 2000;21:32-3.
5. WHO. The Reproductive Health of Adolescents: A Strategy for Action, A joint WHO/UNFPA/ UNICEF Statement. Geneva: WHO; 1989.
6. International Institute for Population Science. National Family Health Survey NFHS 3 India. Mumbai, India: International Institute for Population Science; 2005-06.
7. NIHFWS. Adolescent Health. Reproductive and Child Health for Medical Officers. New Delhi: NIHFWS; 2001. p. 662-5.
8. Shanbhag D, Shilpa R, D'Souza N, Josephine P, Singh J, Goud BR. Perceptions regarding menstruation and Practices during menstrual cycles among high school going adolescent girls in resource limited settings around Bangalore city, Karnataka, India. *Int J Collab Res Intern Med Public Health* 2012;7:1353-62.
9. Sachan B, Idris MZ, Jain S, Kumari R, Singh A. Nutrient intake among school going adolescent girls in Lucknow District. *J Biomed Pharma Res* 2013;2:20-5.
10. Baliga SS, Naik AV, Mallapur DM. Nutritional status of adolescent girls residing in rural area. *J Sci Soc* 2014;41:22-5.
11. Guduri BG, Avvaru K, Naidu AS. To evaluate the prevalence of TMJ disorders among the group of school going adolescents who are undergoing orthodontic treatment. *IOSR J Dent Med Sci* 2014;31:31-3.
12. Yerpude PN, Jogdand KS, Jogdand M. A study of health status among school going adolescent girls in South India. *Int J Health Sci Res* 2013;3:8-12.
13. Sachan B, Idris MZ, Jain S, Kumari R, Singh A. Social determinants and its influence on the prevalence of morbidity among adolescent girls. *N Am J Med Sci* 2012;4:474-47.

**How to cite this article:** Arya V, Chaturvedi S. Comparative study on morbidity pattern among adolescent schoolgirls in rural and urban school of Jhansi. *Int J Med Sci Public Health* 2018;7(5):345-348.

**Source of Support:** Nil, **Conflict of Interest:** None declared.