

# Effectiveness of interventions on biological monitoring among workers exposed to Pb from lead-acid storage battery plant

Ravibabu Kalahasthi, Tapu Barman, Rajmohan HR, Bhavani Shankara Bagepally, Ravichandran Beerappa

Regional Occupational Health Centre (Southern), Indian Council of Medical Research, Kannamangala (Post), Devanahalli, Karnataka, India.  
Correspondence to: Ravibabu Kalahasthi, E-mail: kalahasthi20012002@yahoo.co.in

Received November 19, 2015. Accepted January 5, 2016

## Abstract

**Background:** Studies related to the effectiveness of combined interventions such as engineering and administrative controls, use of a respirator, and conducting awareness program about lead exposure on biological monitoring among lead battery workers were not reported.

**Objective:** This study has assessed the effectiveness of interventions on biological monitoring among lead battery manufacturing workers.

**Materials and Methods:** The intervention methods were use of a respirator, awareness program about lead exposure, engineering and administrative controls. Blood lead levels (BLLs) among workers were estimated using an atomic absorption spectrophotometer. Hematological parameters in workers were determined using Fx-19E automated hematology analyzer. Reticulocyte count in workers was done using supravital staining method.

**Results:** The mean systolic and diastolic blood pressure, the proportion of alcohol consumption, and smoking habits were noted to be significantly decreased among workers after the intervention. The BLLs were reduced significantly among workers engaged in assembly, casting, engineering services, formation, acid filling, and in executives after the intervention. The red cell indices such as erythrocytes and hematocrit were significantly increased and mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, and reticulocyte count were significantly decreased after the intervention.


**Conclusions:** Austere implementation of control measures in lead-acid storage battery plant showed beneficial health effects among lead exposed workers.

**KEY WORDS:** Blood lead levels, interventions, biological monitoring, lead battery workers

## Introduction

Manufacturing of lead-acid storage batteries involves several processes, such as preparation of lead oxide, grid casting, pasting, plate cutting, formation, charging, and assembly.

The chemicals used in these processes are of hazardous nature, and these include lead oxide (PbO<sub>2</sub>), spongy lead (Pb), and sulfuric acid (H<sub>2</sub>SO<sub>4</sub>). Lead enters the body of workers by inhalation and ingestion. After entry into the body, the lead will accumulate in erythrocytes, soft tissue (brain, kidney, and bone marrow), and mineralized tissue (bone, teeth).<sup>[1]</sup> Chronic exposure to Pb during the lead battery manufacturing process leads to a variety of health problems. Previous studies have reported association between blood lead levels (BLLs) with neurobehavioral impairments,<sup>[2]</sup> genotoxic effects,<sup>[3]</sup> impaired coagulation,<sup>[4]</sup> health illness,<sup>[5]</sup> hypertension,<sup>[6]</sup> decreased δ-aminolevulinic acid dehydratase & elevated δ-aminolevulinic acid,<sup>[7]</sup> ocular changes,<sup>[8]</sup> respiratory, gastrointestinal & musculo-skeletal morbidities,<sup>[9]</sup> epigenetic changes,<sup>[10]</sup> decreased bone mineral density,<sup>[11]</sup> oxidative stress,<sup>[12]</sup> dental problems,<sup>[13]</sup> altered

Access this article online	
Website: <a href="http://www.ijmsph.com">http://www.ijmsph.com</a>	Quick Response Code: 
DOI: 10.5455/ijmsph.2016.19112015318	

International Journal of Medical Science and Public Health Online 2016. © 2016 Ravibabu Kalahasthi. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material for any purpose, even commercially, provided the original work is properly cited and states its license.

gene polymorphisms,<sup>[14–16]</sup> increased risk of cardiovascular disease,<sup>[17]</sup> anemia & abdominal pain<sup>[18]</sup>; and decreased enzymatic antioxidants<sup>[19]</sup> among lead battery manufacturing workers.

Sato and Yano<sup>[20]</sup> reported hand contamination was positively associated with BLLs. The variety of household interventions such as home wet cleaning activity; dust control; parental knowledge about lead exposure; and reduction of lead in soil, petrol, and paints will reduce lead poisoning in children.<sup>[21–23]</sup> Dyosi<sup>[24]</sup> studied the effect of interventions of engineering and administrative controls and use of personal protective equipment for lead exposure from lead smelting process and suggested that engineering controls were rated the best control measure. Lee *et al.*<sup>[25]</sup> assessed the use of respirators on biological monitoring among lead battery workers. The result of the study found a significant decrease of BLLs,  $\delta$ -aminolevulinic acid, and lead-related symptoms. Studies related to effectiveness of combined interventions such as engineering and administrative controls, use of a respirator, and conducting awareness program about lead exposure on the status of biological monitoring among workers from lead-acid storage battery manufacturing plant were not reported. The present study was carried to assess the effectiveness of interventions on biological monitoring among workers involved in lead-acid storage battery plant.

## Materials and Methods

The study design was quasi-experimental time series, the data were collected in two phases with pre- and postintervention. The preintervention phase assessment was conducted during July 2013 and postintervention assessment was conducted during July 2014. Informed consent was taken from each worker before participation in the study. The study was approved by Institutional Ethical Committee. In pre- and postintervention phases, the demographic details, blood pressure, BLLs, and hematological parameters were assessed in 397 male lead workers involved in lead-acid storage battery plant.

### Intervention Methods

1. Engineering controls: (1) Plate cutting area was relocated and rerouting of exhaust ducts carried out to improve the assembly work environment. (2) The placing of roof-mounted humidifiers to provide cold air for worker comfort and dust suppression. (3) Installation of the water bath at plate cutting area to trap dust emissions.

2. Administrative controls: (1) Motorized road cleaning machine was introduced to minimize the dust exposure. (2) De-lead hand wash solution was used to avoid hand contamination in workers.

3. Personal protective equipment: Workers used half face mask (V 800) for respiratory protection during their work.

4. An awareness program about lead exposure was conducted.

### Demographic Details

The demographic details such as age, sex, experience, body mass index (BMI), occupational health history, smoking, and alcohol consumption habits among workers were collected using a predesigned validated questionnaire. BMI was calculated by noting the height and weight of each worker and expressed as kg/m<sup>2</sup>.

### Blood Pressure

A health physician measured the blood pressure of on right arm, while the worker was in a sitting position. A standard mercury sphygmomanometer worker was used for blood pressure monitoring.

### Blood Lead Level

Three milliliters of venous whole blood was collected in heparinized vacuette from the workers and stored at –20°C until analysis. Two milliliters of blood was digested by using an ETHOS-D, milestone microwave system (Italy) with 2 mL of nitric acid (HNO<sub>3</sub>) and 0.2 mL of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by maintaining power, temperature, duration of time. The digested samples were made up to 5 mL using triple distilled water and centrifuged. The concentration of lead in blood was measured by using an atomic absorption spectrophotometer (GBC-Avanta P. Australia). A standard solution of 20 µg/dL was prepared from the lead standard solution of Merck (1.19776.0500) and added to the lowest concentration of the sample and the analysis found 100% of recovery with % RSD at <0.5 for three replicates. BLLs were expressed as µg/dL.

### Hematological Parameters

A volume of 2 mL of venous whole blood was collected in a Lavender Top Vacutainer Tube, which contains tripotassium–EDTA. The levels of red blood cell (RBC) count, hematocrit mean corpuscular volume, mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) were determined by a Fx-19E automated hematology analyzer.

### Reticulocyte Count

Non-nucleated immature RBCs contain nuclear remnants of RNA known as a reticulocyte. The reticulocyte counts in peripheral blood samples were done using the supra-vital staining method. The reticulocyte counts in the workers were expressed as percentage.

### Statistical Analysis

Software package SPSS, version 16.0 for windows was used for the analysis of data. The data of age, experience, BMI, blood pressure (BP) (systolic BP [SBP] and diastolic BP [DBP]), BLLs, and hematological parameters were presented as mean ± standard deviation. The effectiveness of interventions on BLLs was assessed using McNemar test. Smoking and alcohol consumption habits among workers were reported in percentage and compared using the chi square test. The probability value of less than 0.05 considered as significant.

## Results

The demographic details among workers in pre- and postintervention groups were presented in Table 1. The mean SBP DBP the proportion of smoking and alcohol consumption habits were significantly decreased in the postintervention group as compared to the preintervention group. No significant difference was observed with respect to age, experience and BMI.

BLLs among workers, according to their job categories of pre- and postintervention groups, were reported in Table 2. The BLLs were significantly decreased in workers from assembly, casting, engineering services, formation, acid filling in executives. No significant differences were observed in pasting, ball mill, charging, and plate cutting area.

BLLs between pre- and postintervention groups according to Biological Exposure Index of American Conference of Governmental Industrial Hygienist was presented in Figure 1. The effectiveness of the intervention on BLLs was assessed using the McNemar test, which showed significantly ( $P < 0.001$ ) decreased BLLs.

Hematological parameters among workers, according to pre- and postintervention groups were reported in Table 3. The levels of erythrocyte count and hematocrit were significantly increased and MCH, MCHC, and reticulocyte count were significantly decreased in the postintervention group as compared to the preintervention group.

## Discussion

The present study assessed the effectiveness of interventions on biological monitoring among workers exposed to lead from lead-acid-storage battery plant. The intervention methods adopted were engineering administrative controls, use of respirators, and awareness program conducted about

**Table 1:** Demographic detail of workers in pre- and postintervention groups

Variables	Preintervention (n = 397)	Postintervention (n = 397)
Age (years)	36.0 ± 4.8	37.0 ± 4.9
Experience (years)	12.6 ± 3.9	13.8 ± 4.1
BMI (kg/m <sup>2</sup> )	25.6 ± 2.9	25.6 ± 2.8
SBP (mm/Hg)	128 ± 14	123 ± 13**
DBP (mm/Hg)	78 ± 11	73 ± 10**
Smoking habits (%)	93 (23)	77 (19)**
Alcohol consumption	213 (54)	203 (51)*

\*\* $P < 0.001$  and \* $P < 0.05$ .

lead exposure. During the study, SBP, DBP, proportion of alcohol consumption, and smoking habits were significantly reduced in the postintervention group as compared to the preintervention group. Dickenson *et al.*<sup>[26]</sup> reported that lifestyle intervention such as weight reduction and restriction of alcohol, smoking, and salt intake significantly reduced the BP as compared to the nonintervention group. The altered lifestyle factors might have influenced the reduction of SBP and DBP in the postintervention group. A recent study reported that BLLs were associated with elevated BP.<sup>[27]</sup> The reduction in BLLs might have influenced the reduction BP among workers. Karita *et al.*<sup>[28]</sup> reported that the nonsmokers have decreased BLLs as compared to smokers. In the present study, the decreased proportion of smoking habits among postintervention might contribute to the reduction of BLLs. This finding was similar to that of Chuang *et al.*<sup>[29]</sup> as restriction of smoking and eating at work places influences reduction in BLLs.

Ochieng *et al.*<sup>[30]</sup> studied the effectiveness of health education and hygiene practices on BLLs among lead battery workers in Nairobi. The authors concluded that use of Personal Protective Equipment (PPE) and health education significantly associated with decreased BLLs. Lormphongs


**Table 2:** Blood lead levels ( $\mu\text{g}/\text{dL}$ ) among workers according to job categories in pre- and postintervention groups

Job categories	Preintervention	Postintervention	Difference in means	Decrease (%)
Assembly (n = 193)	34.6 ± 12.8	28.0 ± 11.4**	6.6	19.0
Casting (n = 40)	29.3 ± 9.8	23.8 ± 10.7*	5.5	18.7
Pasting (n = 37)	33.5 ± 12.1	33.2 ± 13.1	0.3	0.9
Ball mill (n = 9)	34.3 ± 7.7	27.2 ± 11.0	7.1	20.7
Charging (n = 18)	22.3 ± 7.0	17.2 ± 9.3	5.1	22.8
Plate cutting (n = 6)	37.0 ± 14.6	31.3 ± 7.4	5.7	15.4
Engineer services (n = 48)	31.2 ± 12.0	22.1 ± 9.4**	9.1	29.2
Formation (n = 4)	26.5 ± 6.5	13.5 ± 2.9*	13.0	49.5
Acid filling (n = 10)	29.8 ± 9.5	21.6 ± 9.7*	8.2	29.5
Executives (n = 32)	35.6 ± 14.0	27.1 ± 12.4*	8.5	23.8
Total	32.8 ± 12.3	26.5 ± 11.7**	6.3	19.2

\*\* $P < 0.001$  and \* $P < 0.05$ .

**Table 3:** Hematological parameters in workers according to pre- and postintervention groups

Variables	Preintervention	Postintervention
RBC (million/ $\mu$ L)	4.23 $\pm$ 0.38	4.32 $\pm$ 0.46*
Hematocrit (%)	36.9 $\pm$ 3.0	37.7 $\pm$ 4.2*
MCV (fl)	87.7 $\pm$ 7.1	87.6 $\pm$ 7.8
MCH (pg)	34.5 $\pm$ 3.3	32.6 $\pm$ 4.6*
MCHC (%)	39.3 $\pm$ 2.3	37.2 $\pm$ 3.8*
Reticulocyte count (%)	0.84 $\pm$ 0.3	0.66 $\pm$ 0.2*

\* $P < 0.05$ .**Figure 1:** Blood leads levels among pre- and postintervention workers according to Biological Exposure Index—ACGIH criteria.

*et al.*<sup>[31]</sup> also reported that the occupational health education causes decreased BLLs among lead-acid-storage battery workers. Esswein *et al.*<sup>[32]</sup> reported that hand contamination was positively associated with BLLs. Dyosi<sup>[24]</sup> studied the effect of engineering and administrative controls and use of personal protective equipment for lead exposure from lead smelting process and suggested engineering controls were rated the best control measure. Lee *et al.*<sup>[25]</sup> assessed the effectiveness of the respirator on biological monitoring on the health management of lead battery workers. That study found significantly decreased BLLs,  $\delta$ -aminolevulinic acid, and lead-related symptoms. In the present study, engineering administrative controls and use of respirators were implemented and also awareness program was conducted to minimize lead exposure. BLLs were significantly reduced in assembly, casting, engineering services, formation, acid filling and in executive job categories of postintervention group. The present study also assessed the hematological parameters in pre- and postintervention groups. The mean levels of erythrocytes and hematocrit significantly increased and MCH, MCHC, and reticulocyte counts were significantly

decreased in the postintervention group as compared to the preintervention group. Saichanma *et al.*<sup>[33]</sup> also noticed higher level of the RBC count with low levels of MCH and MCHC. The findings of this study were similar to previous studies.

## Conclusion

The variables of BLLs, hematological parameters, and lifestyle factors were significantly improved in the post-intervention as compared to the preintervention group. These study findings indicate the beneficiary effects in terms of biological monitoring with the austere implementation of control measure in the lead-acid battery plant.

## References

1. WHO/ Lead poisoning and Health. (2014), Available from: <http://www.who.int/mediacentre/factsheets/fs379/en/> pages: 1-4.
2. Ravibabu K, Barman T, Rajmohan HR. Serum neuron-specific enolase, biogenic amino-acids and neurobehavioral function in lead-exposed workers from lead-acid battery manufacturing process. *Int J Occup Environ Med* 2015;6(1):50–7.
3. Chinde S, Kumari M, Devi KR, Murty US, Rahman MF, Kumari SI, *et al.* Assessment of genotoxic effects of lead in occupationally exposed workers. *Environ Sci Pollut Res Int* 2014;21(19):11469–80.
4. Barman T, Kalahasthi R, Rajmohan HR. Effects of lead exposure on the status of platelet indices in workers involved in a lead-acid battery manufacturing plant. *J Expo Sci Environ Epidemiol* 2014;24(6):629–33.
5. Ahmad SA, Khan MH, Khandker S, Sarwar AF, Yasmin N, Faruquee MH, *et al.* Blood lead levels and health problems of lead acid battery workers in Bangladesh. *Scientific World Journal* 2014;2014:974104.
6. Were FH, Moturi MC, Gottesfeld P, Wafula GA, Kamau GN, Shiundu PM. Lead exposure and blood pressure among workers in diverse industrial plants in Kenya. *J Occup Environ Hyg* 2014;11(11):706–15.
7. Petracca M, Scafa F, Boeri R, Flachi D, Candura SM. Imported occupational lead poisoning: report of four cases. *Med Lav* 2013; 104(6):428–33.
8. Ekinci M, Ceylan E, Cagatay HH, Keleş S, Altinkaynak H, Kartal B, *et al.* Occupational exposure to lead decreases macular, chorioidal, and retinal nerve fiber layer thickness in industrial battery workers. *Curr Eye Res* 2014;39(8):853–8.
9. Kalahasthi RB, Barman T, Rajmohan HR. The relationship between blood lead levels and morbidities among workers employed in a factory manufacturing lead-acid storage battery. *Int J Environ Health Res* 2014;24(3):246–55.
10. Li C, Yang X, Xu M, Zhang J, Sun N. Epigenetic marker (LINE-1 promoter) methylation level was associated with occupational lead exposure. *Clin Toxicol (Phila)* 2013;51(4):225–9.
11. Raafat BM, Hassan NS, Aziz SW. Bone mineral density (BMD) and osteoporosis risk factor in Egyptian male and female battery manufacturing workers. *Toxicol Ind Health* 2012;28(3):245–52.
12. Conterato GM, Bulcao RP, Sobieski R, Moro AM, Charao MF, de Freitas FA, *et al.* Blood thioredoxin reductase activity, oxidative stress and hematological parameters in painters and battery workers: relationship with lead and cadmium levels in blood. *J Appl Toxicol* 2013;33(2):142–50.

13. Suyama Y, Takaku S, Okawa Y, Matsukubo T. Dental erosion in workers exposed to sulfuric acid in lead storage battery manufacturing facility. *Bull Tokyo Dent Coll* 2010;51(2):77–83.
14. Chen HI, Chiu YW, Hsu YK, Li WF, Chen YC, Chuang HY. The association of metallothionein-4 gene polymorphism and renal function in long-term lead-exposed workers. *Biol Trace Elem Res* 2010;137(1):55–62.
15. Shaik AP, Jamil K. Polymorphisms in MGP gene and their association with lead toxicity. *Toxicol Mech Methods* 2009;19(3):209–13.
16. Shaik AP, Jamil K. A study on the ALAD gene polymorphisms associated with lead exposure. *Toxicol Ind Health* 2008;24(7):501–6.
17. Kim KR, Lee SW, Paik NW, Choi K. Low-level lead exposure among South Korean lead workers, and estimates of associated risk of cardiovascular diseases. *J Occup Environ Hyg* 2008;5(6):399–416.
18. Fonte R, Agosti A, Scafa F, Candura SM. Anaemia and abdominal pain due to occupational lead poisoning. *Haematologica* 2007;92(2):e13–4.
19. Patil AJ, Bhagwat VR, Patil JA, Dongre NN, Ambekar JG, Jailkhani R, et al. Effect of lead (Pb) exposure on the activity of superoxide dismutase and catalase in battery manufacturing workers (BMW) of western Maharashtra (India) with reference to heme biosynthesis. *Int J Environ Res Public Health* 2006;3(4):329–37.
20. Sato M, Yano E. The association between lead contamination on the hand and blood lead concentration: a workplace application of the sodium sulphide (Na<sub>2</sub>S) test. *Sci Total Environ* 2006;363(1–3):107–13.
21. Yeoh B, Woolfenden S, Lanphear B, Ridley GF, Livingstone N, Jorgensen E. Household interventions for preventing domestic lead exposure in children. *Cochrane Database Syst Rev* 2014;12:CD006047.
22. Kennedy C, Lordo R, Sucusky MS, Boehm R, Brown MJ. Primary prevention of lead poisoning in children: a cross-sectional study to evaluate state specific lead-based paint risk reduction laws in preventing lead poisoning in children. *Environ Health* 2014;13:93.
23. Mielke HW, Covington TP, Mielke PW Jr, Wolman FJ, Powell ET, Gonzales CR. Soil intervention as a strategy for lead exposure prevention: the New Orleans lead-safe childcare playground project. *Environ Pollut* 2011;159(8–9):2071–7.
24. Dyosi S. Evaluation of preventive and control measures for lead exposure in a South African lead-acid battery recycling smelter. *J Occup Environ Hyg* 2007; 4(10):762–9.
25. Lee BK, Lee CW, Ahn KD. The effect of respiratory protection with biological monitoring on the health management of lead workers in a storage battery industry. *Int Arch Occup Environ Health* 1993;65(Suppl 1):S181–4.
26. Dickinson HO, Mason JM, Nicolson DJ, Campbell F, Beyer FR, Cook JV, et al. Lifestyle interventions to reduce raised blood pressure: a systematic review of randomized controlled trials. *J Hypertens* 2006;24(2):215–33.
27. Vupputuri S, He J, Muntner P, Bazzano LA, Whelton PK, Batuman V. Blood lead level is associated with elevated blood pressure in blacks. *Hypertension* 2003;41:463–8.
28. Karita K, Nakao M, Awake K, Yamanouchi Y, Nishikitani M, Nomura K, et al. Blood lead and erythrocyte protoporphyrin levels in association with smoking and personal hygienic behavior among lead exposed workers. *Occup Environ Med* 2005;62:300–3.
29. Chuang HY, Lee ML, Chao KY, Wang JD, Hu H. Relationship of blood lead levels to personal hygiene habits in lead battery workers: Taiwan, 1991–1997. *Am J Ind Med* 1999;35(6):595–603.
30. Ochieng NJ. Effect of Health Education and Hygiene on Blood Lead Levels, Among Lead Battery Workers in Nairobi and Athi River, Kenya. Thesis of MPH, School of Public Health, Kenyatta University, 2012.
31. Lormphongs S, Morioka I, Miyai N, Yamamoto H, Chaikittiporn C, Thiramanus T, et al. Occupational health education and collaboration for reducing the risk of lead poisoning of workers in a battery manufacturing plant in Thailand. *Ind Health* 2004;42:440–5.
32. Esswein EJ, Boeniger MF, Ashley K. Handwipe method for removing lead from skin. *J ASTM Intl* 2011;8(5). (DOI: 10.1520/JAI103527)
33. Saichanma S, Chulsomlee S, Thangrua N, Pongsuchart P, Sanmun D. The observation report of red blood cell morphology in Thailand teenager by using data mining technique. *Adv Hematol* 2014;2014:493706.

**How to cite this article:** Kalahasthi R, Barman T, Rajmohan HR, Bagepally BS, Beerappa R. Effectiveness of interventions on biological monitoring among workers exposed to Pb from lead-acid storage battery plant. *Int J Med Sci Public Health* 2016;5:1770-1774

**Source of Support:** Nil, **Conflict of Interest:** None declared.