

# PREVALENCE OF OVERWEIGHT AND OBESITY AND ITS ASSOCIATION WITH HYPERTENSION AND DIABETES: A STUDY AMONG MANAKKARAMBAI POPULATION OF THANJAVUR DISTRICT

C Manoji Jayasingh<sup>1</sup>, A Sheeba Christina<sup>2</sup>

<sup>1</sup> Department of Biochemistry, Vinayaka Missions Medical College, Karaikal, Pondicherry, India

<sup>2</sup> Department of Biotechnology, PRIST University, Thanjavur, Tamil Nadu, India

Correspondence to: A Sheeba Christina (sheb\_chris@yahoo.com)

DOI: 10.5455/ijmsph.2013.200420132

Received Date: 20.03.2013

Accepted Date: 20.04.2013

## ABSTRACT

**Background:** Obesity is a well-known risk factor of hypertension and diabetes.

**Aims & Objective:** This study was aimed to investigate the prevalence of overweight and obesity in Manakkarambai population and to evaluate its relationship with hypertension and diabetes.

**Material and Methods:** This Cross sectional study comprises of 163 subjects attending a local medical camp between 25- 70 years of age. Height, weight, BMI, systolic and diastolic blood pressure, fasting blood sugar and post prandial blood sugar were determined using standard methods.

**Results:** Prevalence of overweight and obesity was 15% and 34% respectively and it was predominant among women than men. Systolic blood pressure, diastolic blood pressure and post prandial blood sugar were significantly elevated in obese subjects than in normal BMI subjects (control).

**Conclusion:** The results showed a higher trend of hypertension and diabetes with increase in BMI.

**KEY-WORDS:** Overweight; Obesity; Hypertension; Diabetes; Manakkarambai

## Introduction

The issue of overweight and obesity has become a serious public health concern throughout the world in the past few decades. In India, it has reached epidemic proportion in the 21<sup>st</sup> century with morbid obesity affecting 5% of the country's population.<sup>[1]</sup> Although the terms overweight and obesity are often used interchangeably, overweight refers to an excess of body weight compared with height; obesity refers to an excess of body fat. BMI is the most widely used measure to define obesity. It is determined by dividing weight (wt) in kilograms by height (ht) in meters square.<sup>[2]</sup> A desirable BMI according to the WHO recommended cut-offs for Asians is considered to be between 18.5 and 22.9 kg/m<sup>2</sup>. A BMI of 23–24.9 kg/m<sup>2</sup> is defined as overweight and >25 kg/m<sup>2</sup> as 'obese'.<sup>[3]</sup> A strong association has been depicted between BMI and mortality among Asian populations.<sup>[4]</sup> BMI is also positively and independently associated with morbidity and mortality from hypertension and type II diabetes

mellitus.<sup>[5]</sup> Relationships between obesity and the risk for hypertension and diabetes should be assessed early, if not, it might lead to micro and macrovascular complications. Though the association of BMI with hypertension and diabetes has long been the subject of epidemiological research, it has not been sufficiently explored in rural population. Hence the present study was aimed to find the prevalence of overweight and obesity in this segment of rural population and to explore the relation of BMI with hypertension and diabetes.

## Materials and Methods

A cross sectional study was carried out in 163 participants attending a free medical camp at Manakkarambai village, belonging to the age group 25-70. Gender-wise distribution of 163 subjects was 92 males and 71 females. BMI was measured from height and weight using the standard protocol of Weiner and Lourie.<sup>[6]</sup> Height was measured using height measuring scale with

0.1cm division value and weight using potable weighing machine without shoes. Blood pressure was determined using the standard mercury sphygmomanometer. Normal BP is defined as less than 120/80 mmHg, pre-hypertension as 120-139/80-89 mmHg, stage I hypertension as 140-159/ 90-99 mmHg, and stage II hypertension as equal to and above 160/100 mmHg.<sup>[7]</sup> Fasting and post prandial blood sugar was determined using glucometer. Written informed consent was obtained from all subjects.

### Statistical Analysis

Statistical analysis of data was done using SPSS (version 14). Values are expressed as mean value. Standard deviation was done to obtain accuracy. One way ANOVA was performed to determine the significance and P value less than 0.05 was considered significant.

## Results

Total participants who attended the free medical camp were divided into 4 groups based on BMI namely, normal, underweight, overweight and obese. Figure 1 shows the prevalence of overweight, obesity, hypertension and diabetes in Manakkarambai population. Among 163 participants screened, overweight and obese subjects correspond to 15% and 34% respectively. It was found to be predominant among women (9 and 19%) than men (6 and 15%). Occurrence of underweight subjects was 16%. Prevalence of hypertension was 20% comprising of 13% of men and 7% women. In case of diabetes, 5% of men and 6% of women put up to a total of 11% of diabetic subjects in this segment of population.

The mean values of various parameters in different categories of BMI were given in Table 1. In case of obese subjects, Systolic BP (136.60 mmHg), Diastolic BP (86.96 mmHg) and post-prandial blood sugar (160.59mg/dl) were significantly elevated than in that of normal BMI subjects (122.45 mmHg, 78.07 mmHg and 134.26 mg/dl). In case of overweight and subjects, significant increase was observed only in case of post-prandial blood sugar level (141.12 mg/dl) when compared with the normal-weight subjects

(134.26 mg/dl) whereas in the other parameters like systolic and diastolic BP and fasting blood sugar the difference was not found to be significant. No significant difference was seen in all the parameters between under-weight and normal weight subjects respectively.

Table 2 gives the distribution of study subjects in different classes of BMI and BP. Among normal weight subjects, 35% have normal BP and 65% are in pre-hypertensive stage. The same trend was also seen in underweight subjects. Out of 26 subjects, 9 (35%) have normal BP and 17 (65%). Thus in both the groups no subject was found to be hypertensive. In overweight subjects, the percentage of subjects with normal BP, pre-hypertension and stage I hypertension was 17%, 71% and 12% respectively. In case of obese subjects, all 4 BP classes were recorded namely, 5% with normal BP; 57% in pre-hypertensive stage; 16% in stage I and 22% in stage II hypertension. Thus stage I and stage II hypertension were seen only in overweight and obese subjects.

Distribution of study subjects in IGT and diabetic condition was recorded in Table 3. Among normal weight subjects, only 2% was observed in diabetic condition. The remaining population namely 79% have normal blood sugar level and 18% in IGT condition. In Underweight subjects 85% have normal blood sugar and only 15% in IGT condition. No diabetic subject was seen in this group. In case of overweight and obese subjects, the percentage of population in normal, IGT and diabetic condition was 33%, 42%, 25% and 36%, 46%, 18% respectively.

**Table-1: Mean values of Various Parameters in Different BMI Classes**

BMI Classes	SBP mm/Hg	DBP mm/Hg	FBS mg/dl	PPBS mg/dl
Normal Weight (n=57)	122.45 (10.90)	78.07 (5.48)	103.24 (17.45)	134.26 (24.09)
Underweight (n=26)	118.8 (9.08)	78.46 (6.12)	97.11 (14.42)	131.5 (14.89)
Overweight (n=24)	125.83 (10.17)	78.33 (6.37)	107.54 (12.97)	141.12 (21.20)*
Obese (n=56)	136.60 (16.65)**	86.96 (10.43)*	105.92 (12.18)	160.59 (39.49)*

SBP: Systolic Blood Pressure; DBP: Diastolic Blood Pressure; FBS: Fasting Blood Sugar; PPBS: Post Prandial Blood Sugar; Values are represented as mean (standard deviation); \* Values are significant at 5% level; \*\* Values are significant at 1% level

**Table-2: Distribution of Subjects in Different Categories of BMI and BP**


BMI Classes	Normal BP	Pre-HT	Stage I HT	Stage II HT
Normal Weight (n=57)	20 (35%)	37 (65%)	0	0
Underweight (n=26)	9 (35%)	17 (65%)	0	0
Overweight (n=24)	4 (17%)	17 (71%)	3 (12%)	0
Obese (n=56)	3 (5%)	32 (57%)	9 (16%)	12 (22%)

Values are expressed as No. of subjects and their percentage in parenthesis; HT: Hypertension

**Table-3: Prevalence of IGT and Diabetes in different categories of BMI**

BMI Classes	Normal Blood Sugar	IGT	Diabetes
Normal Weight (n=57)	45 (79%)	10 (18%)	2 (3%)
Underweight (n=26)	22 (85%)	4 (15%)	0
Overweight (n=24)	8 (33%)	10 (42%)	6 (25%)
Obese (n=56)	20 (36%)	26 (46%)	10 (18%)

Values are expressed as No. of subjects and their percentage in parenthesis

**Figure-1: Gender-wise Occurrence of Overweight and Obesity and Its Associated Disorders in Manakkarambai Population**

## Discussion

We observed a fairly high prevalence of overweight and obesity among Manakkarambai population. Among 163 participants screened, overweight and obese subjects correspond to 15% and 34% respectively. Prevalence of obesity was found to be quite high in this rural community as it is almost in par with the normal population (35%). In the past, the overweight prevalence was traditionally higher in urban areas. However, the opposite trend applies nowadays. This could be because subjects living in rural areas have poorer access to health care and they practice medical self-care less.<sup>[8]</sup> Increase in body weight may also be due to the change in lifestyle factors which, has also started affecting the rural community like fat-rich diet, more usage of alcohol and tobacco,

increased energy intake and low energy output due to lesser involvement in physical activities.

Based on gender, overweight and obesity were predominant among female (9% and 19%) that in male (6% and 15%). Similar observations have been reported in India by Gothankar.<sup>[1]</sup> Worldwide, gender differences are more pronounced, with 10% of men and 14% of women obese.<sup>[9]</sup> Similarly, the prevalence of hypertension and diabetes were 20% and 11% respectively. Hypertension was more pronounced among male (13%) than in female (7%), which is consistent with the findings of Humayun et al.<sup>[10]</sup> whereas much difference was not observed in diabetic subjects of both gender. However in obese subjects, Bhat et al.<sup>[11]</sup> has reported high prevalence of hypertension in female than in male.

The relationship between BMI and hypertension is of particular interest to developing countries as excess cardiovascular mortality among lean hypertensive subjects has been reported in some longitudinal studies.<sup>[12]</sup> The association between hypertension and BMI is evident from table 2. In normal and underweight subjects majority of them are in the pre-hypertension stage (65%). In overweight subjects, the % of population in pre-hypertension stage is more (71%) and also 12% in stage I hypertension. Similarly in obese population 16% in stage I and 22% in stage II hypertension respectively. The important point is that only obese patients are in stage II hypertension. This clearly indicates the positive association of hypertension with BMI. This is consistent with the findings of Mungreiphy et al.<sup>[4]</sup>, Humayun et al.<sup>[10]</sup> and Ahmed et al.<sup>[13]</sup>.

BMI is an indicator of obesity and would be a possible indicator of the risk of diabetes. Prevalence of diabetes in overweight and obese subjects was 25% and 18% respectively, whereas in normal population it is only 3% (Table 3). Diabetes prevalence was significantly higher in overweight and obese subjects than in normal-weight subjects. Among overweight and obese subjects 42% and 46% were in IGT condition. Gothankar<sup>[1]</sup>, have also reported a positive association of BMI with diabetes. BMI is an indicator of obesity and would be a possible indicator of the risk of Diabetes Mellitus.<sup>[9]</sup>

## Conclusion

Overweight and obesity has become a major health problem in this rural community. The study also indicates a positive association of BMI with hypertension and diabetes. Significant number of subjects are seen in pre-hypertension and IGT condition which clearly indicates and the need of creating awareness regarding overweight and obesity and related chronic diseases like hypertension and diabetes which would otherwise become emerging problems in the near future.

## References

1. Gothankar JS. Prevalence of obesity and its associated comorbidities amongst adults. *National Journal of Community Medicine*. 2011; 2(2):221-224.
2. Brown CD, Higgins M, Donato KA, Rohde FC, Garrison R, Obarzanek E, et al. Body Mass Index and the Prevalence of Hypertension and Dyslipidemia. *Obes Res*. 2000;8:605-619.
3. Low S, Chin MC, Ma S, Heng DM, Deurenberg Y. Rationale for Redefining Obesity in Asians. *Ann Acad Med Singapore*. 2009;38:66-74.
4. Mungreiphy NK, Kapoor S, Sinha R. Association between BMI, Blood Pressure, and Age: Study among Tangkhul Naga Tribal Males of Northeast India. *Journal of Anthropology*. 2011;2011:6.
5. Pi-Sunyer FX. Medical hazards of obesity. *Annals of Internal Medicine*. 1993 ;119(7 Pt 2):655-60.
6. Weiner JS, Lourie JA. *Practical Human Biology*. Academic Press, London, UK, 1981.
7. Braunwald E, Fauci AS, Kasper DL, Hauser SL, Longo DL, Jameson JL. *Harrison's Principle of Internal Medicine*. Vol.1. 15<sup>th</sup> ed. New York: Tata McGraw- Hill publishers. 2001. Pp- 571-578.
8. Hernandez-Mijares A, Sola-Izquierdo E, Ballester-Mecho F, Mari-Herrero MT, Gilabert-Moles JV, Gimeno-Clemente N, et al. Obesity and overweight prevalences in rural and urban populations in East Spain and its association with undiagnosed hypertension and Diabetes Mellitus: a cross-sectional population-based survey. *BMC Res Notes*. 2009; 2 :151-154.
9. World Health Organization. *Global Health Observatory*. *Obes Res*. 2002, 10(5):345-50.
10. Humayun A, Shah AS, Sultana R. Relation of hypertension with BMI and age in male and female population of Peshwar, Pakistan. *J Ayub Med Coll Abbottabad*. 2009, 21(3): 63-65.
11. Bhat NA, Kamili MA, Allaqaband GQ. Hypertension in South Kashmir. *Indian Pract*. 2002, 55(4):215-218.
12. Tesfaye F, Nawi NG, Minh HV, Bypass P, Berhane Y, Bonita R, et al. Association between BMI and BP across 3 populations in Africa and Asia. *J Human Hypertension*. 2007, 21: 28- 37.
13. Ahmad R, Khan TM, Zulfiqar S, Marwat MA, Rehman I. Role of Body Mass Index (BMI) in the development of hypertension in adult population of District Swat. *Biomedica*. 2007, 23:39-41.

**Cite this article as:** Jayasingh CMS, Christina AS. Prevalence of overweight and obesity and its association with Hypertension and Diabetes: A study among Manakkarambai population of Thanjavur district. *Int J Med Sci Public Health* 2013; 2:618-621.

**Source of Support: Nil**

**Conflict of interest: None declared**