

# Study of effects of yoga and pranayam on human reaction time and certain physiological parameters in normal and hypertensive subjects

Tejaswini D Sonwane<sup>1</sup>, Neelam V Mishra<sup>2</sup>

<sup>1</sup>Department of Physiology, Seth GS Medical College and KEM Hospital, Mumbai, Maharashtra, India.

<sup>2</sup>Department of Physiology, Government Medical College, Nagpur, Maharashtra, India.

Correspondence to: Tejaswini D Sonwane, E-mail: tejas9444@gmail.com

Received February 4, 2016. Accepted April 1, 2016


## ABSTRACT

**Background:** Yoga is a process of gaining control over the mind, as defined by Patanjali. Stress has been implicated as one of the major causes of essential hypertension. Yoga works on every cell of the body. Yoga influences body as well as controls the stress in the individual. An index of the processing ability of central nervous system and a simple means of determining sensory-motor performance is referred to as reaction time (RT). It has been proclaimed that human performance including central neural processing is improved by yoga training. It improves cardiorespiratory performance, balances autonomic nervous system, decreases pulse, respiratory rate, systolic and diastolic blood pressure. **Aims and Objective:** To show the effects of yoga and pranayam on auditory and visual RT and on certain physiological parameters such as weight, body mass index, pulse rate, respiratory rate, systolic blood pressure, and diastolic blood pressure in normal and hypertensive subjects. **Materials and Methods:** This comparative type of study included 70 normal and 70 hypertensive subjects. It was carried on subjects between 30 and 60 years of age. Auditory reaction time (ART) and visual reaction time (VRT) were studied in subjects with "Response Analyzer" reaction time apparatus. The physiological parameters such as weight, body mass index, pulse rate, respiratory rate, systolic blood pressure, and diastolic blood pressure were measured. Parameters were measured in two sittings; on admission to yoga center and after 3 months. Statistical software STATA Version 10.0 was used for statistical analysis. **Result:** It was found that changes in the RT and physiological parameters were significant in hypertensive subjects when compared with normal individuals. **Conclusion:** Yoga is a helpful intervention in hypertensive subjects. Yoga and pranayam are more beneficial to hypertensive subjects. RT is an index of cortical arousal, and a decrease in it indicates an improved sensory-motor performance and an enhanced processing ability of the central nervous system.

**KEY WORDS:** Yoga; Pranayam; Reaction time; Hypertensive Subject

## INTRODUCTION

Yoga is a process of gaining control over the mind, as defined by Patanjali in his second aphorism (Yoga Sutras: 1.2). Lifestyle-related diseases such as coronary artery disease, obesity, and hypertension are shockingly on the increase in our modern society.<sup>[1-3]</sup> Stress has been implicated as one of the major cause of essential hypertension.<sup>[4-6]</sup> Treatment by weight reduction, yoga, and muscle relaxation each produces smaller but

Access this article online	
Website: <a href="http://www.njppp.com">http://www.njppp.com</a>	Quick Response Code:
DOI: 10.5455/njppp.2016.6.04022016136	

National Journal of Physiology, Pharmacy and Pharmacology Online 2016. © 2016 Tejaswini D Sonwane. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material for any purpose, even commercially, provided the original work is properly cited and states its license.

appreciable changes in blood pressure.<sup>[7-9]</sup> Yogic techniques especially Dhyana and Shavasana improve attentiveness. Increased attentiveness decreases reaction time.

The time interval between the application of a stimulus and the emergence of appropriate voluntary response by a subject is called RT. It involves stimulus processing, decision making, attention mechanism, and response programming.<sup>[10]</sup> With this point of view, we aim to study the degree of effect that yoga brings in normal subjects when compared with hypertensive subjects in terms of human RT, blood pressure, and various other physiological parameters.

This study shows the effects of yoga and pranayam on auditory and visual reaction time and on certain physiological parameters such as weight, body mass index, pulse rate, respiratory rate, systolic blood pressure, and diastolic blood pressure in normal and hypertensive subjects.

### MATERIALS AND METHODS

This comparative type of study was carried out in the yoga center in central India on normal and hypertensive individuals for 3 months. Approval from the Institutional Ethics Committee was obtained. The study included 70 normal and 70 hypertensive

subjects. Planned study population included control group fulfilling the following criteria: Normal healthy adult volunteers between 30 and 60 years of age of both genders, non-smoker, nonalcoholic, not doing yoga previously, and not on any medication. Inclusion criteria for the study group: Subjects with essential hypertension, which included subjects between 30 and 60 years of age with hypertension diagnosed within last 5 years and are on same medication since 5 years, nonsmoker, nonalcoholic, and not doing yoga previously. Exclusion criteria: Subjects with stage-1 hypertension, not on any medication; with stage-2 hypertension; with hypertension with cardiovascular or cerebral complications; with secondary hypertension; and with chronic diseases other than hypertension were excluded from the study. Subjects already practicing yoga and pregnant women were also excluded.

**Planned Intervention:** After taking consent, a detailed history of the subjects was taken. General and systemic examination of the subjects was done. After screening with physical examination, subjects underwent yogic session daily for one-and-a-half hour duration from 6 am to 7.30 am in the morning for a period of 90 days (i.e., 3 months). The session started with *Awayawa-dhyana* and a short prayer, followed by various asanas such as *paschimottanasan*, *vrukshasan*, *tadasan*, *padmasan*, *ardhpadmasan*, *pranayam*, and meditation and ended with a

**Table 1: Effect of yoga on different parameters**

	1st day	After 3 months	P	Mean change	p
Weight (kg)					
Normal	70.11 ± 5.29	67.17 ± 5.42	<0.0001**	3.04 ± 2.77	0.0030**
Hypertensives	71.39 ± 8.20	67.26 ± 7.44	<0.0001**	4.28 ± 1.53	
BMI (kg/m <sup>2</sup> )					
Normal	26.27 ± 2.71	25.23 ± 2.71	<0.0001**	1.15 ± 1.22	0.0055**
Hypertensives	26.12 ± 3.47	24.62 ± 3.15	<0.0001**	1.54 ± 0.59	
Pulse rate (beats/min)					
Normal	72.7 ± 3.19	69.31 ± 2.85	<0.0001**	3.15 ± 2.64	0.0104*
Hypertensives	76.72 ± 4.63	72.24 ± 3.79	<0.0001**	4.78 ± 2.01	
Respiratory rate (respiration/min)					
Normal	14.42 ± 2.37	11.90 ± 1.51	<0.0001**	3.00 ± 2.33	0.0004**
Hypertensives	17.4 ± 2.04	12.9 ± 1.58	<0.0001**	4.37 ± 1.50	
SBP (mm Hg)					
Normal	116.45 ± 2.68	112.05 ± 1.95	<0.0001**	4.63 ± 1.99	0.0004**
Hypertensives	136.11 ± 7.88	130 ± 7.01	<0.0001**	6.20 ± 3.05	
DBP (mm Hg)					
Normal	76.02 ± 2.95	71.73 ± 2.31	<0.0001**	4.70 ± 1.90	0.0013**
Hypertensives	88.0 ± 4.34	82.25 ± 3.79	<0.0001**	5.83 ± 1.83	
ART (milliseconds)					
Normal	181.05 ± 9.93	166.31 ± 6.50	<0.0001**	15.55 ± 6.01	0.0442*
Hypertensives	184.5 ± 10.18	166.78 ± 8.22	<0.0001**	17.88 ± 6.54	
VRT (milliseconds)					
Normal	196.17 ± 5.28	185.25 ± 4.72	<0.0001**	10.91 ± 4.72	<0.0001**
Hypertensives	192.25 ± 8.33	176.35 ± 6.76	<0.0001**	15.9 ± 5.62	

\*p < 0.05, statistically significant; \*\*p < 0.01.

short prayer. Parameters were measured on admission to yoga center (i.e., on the 1st day and after 3 months). Parameters measured were age (years), height (cm), weight (kg), body mass index (BMI) ( $\text{kg}/\text{m}^2$ ), pulse rate (beats/min), respiratory rate (respiration/min), blood pressure (mm Hg). RT (milliseconds) was measured by "Response analyzer."

**Statistical Analysis:** Clinical parameters were compared between the 1st day and after 3 months in normal and hypertensive subjects by performing paired t-test. Categorical variables were compared by  $\chi^2$ -statistics. All the tests were two-sided.  $p < 0.05$  was considered as statistically significant. Statistical software STATA Version 10.0 was used for data analysis.

## RESULT

Weight, BMI, pulse rate, and respiratory rate were found to be significantly decreased in hypertensive subjects when compared with normal subjects. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) also decreased significantly in hypertensive subjects when compared with normal subjects.

The decrease in ART and VRT was statistically significant in hypertensive subjects when compared with normal subjects.

The subjects were given yoga training including asanas, pranayam, and meditation for 3 months. Physiological parameters such as weight, BMI, pulse rate, respiratory rate, SBP and DBP, ART, and VRT were studied at the beginning of the study and after 3 months.

## DISCUSSION

In accordance to our finding, Khadka et al.<sup>[4]</sup> showed significant decrease in body weight after yoga. Khadka et al.<sup>[4]</sup> and Tundwala et al.<sup>[11]</sup> showed statistically significant reduction in BMI, similar to our results.

The mechanism of the beneficial effect of yoga in the management of obesity cannot be explained by simple excess caloric expenditure because practice of asanas does not bring about increased, rapid large muscle activity and energy generation. However, yoga in the management of obesity is of significance.<sup>[12]</sup>

**Pulse Rate (Beats/Min):** Bharshankar et al.,<sup>[13]</sup> Vijayalakshmi et al.,<sup>[14]</sup> and Khadka et al.<sup>[4]</sup> showed significant reduction of pulse rate after yoga training. These results are similar to our study. Slow breathing as in yogic practice enhances heart rate variability and baroreflex sensitivity by resynchronizing inherent cardiovascular rhythms.<sup>[12]</sup> This is owing to increased vagal modulation.<sup>[15,16]</sup> Yoga-based guided relaxation helped in the reduction of sympathetic activity with reduction in heart rate, skin conductance, oxygen consumption, and increase in breath volume.<sup>[17,18]</sup> Statistically significant reduction in pulse rate after regular practice of yoga is attributed to increased vagal tone. Decreased sympathetic activity reduces catecholamine secretion and also leads to vasodilation leading to improvement in

peripheral circulation. Regular yogic practices reduce basal metabolic rate and resting oxygen consumption.<sup>[12]</sup> All these may be responsible for reduction in resting pulse rate.

**Respiratory Rate (Breaths/Min):** Our results are similar to the findings of Doijad and Surdj<sup>[19]</sup> and Abel et al.<sup>[20]</sup> They found significant decrease in RR. Recitation of the yoga mantras (chanting) slowed respiration to almost exactly six breaths per minute and enhanced heart rate variability and baroreflex sensitivity.<sup>[12]</sup> By following pranayama for few weeks, the bulbopontine complex is adjusted to a new pattern of breathing which is slower than its basal rhythm causing decrease in respiratory rate.<sup>[15,19]</sup> During daily practice of pranayama, the fundamental activity of bulbopontine complex is altered in such a way as to bring down its rhythm by voluntarily extending the phases of inspiration and expiration by stretching to their fullest extents, thus making the lungs to work to their maximal extent to take  $\text{O}_2$  and expire  $\text{CO}_2$  maximally.

**Systolic Blood Pressure (mm Hg):** In accordance to our finding, studies done by Sundar et al.,<sup>[21]</sup> Murugesan et al.,<sup>[22]</sup> Dickinson et al.,<sup>[23]</sup> Schmidt et al.,<sup>[24]</sup> Joseph et al.,<sup>[25]</sup> Alexander et al.,<sup>[26]</sup> Steptoe,<sup>[27]</sup> and Patel and Marmot<sup>[28]</sup> showed significant changes in SBP. Different mechanisms have been postulated to explain the blood pressure-lowering effect of weight loss, a reduction of extracellular volume thus decreasing cardiac output, suppression of the sympathetic nervous system, an improvement in insulin resistance, and a normalization of the aldosterone-renin interrelationship.<sup>[29,30]</sup> The blood pressure-lowering effect of weight loss is most likely a result of an improvement in insulin sensitivity and a decrease in sympathetic nervous system activity and occurs independent of salt restriction.<sup>[16]</sup> The mechanism of yoga-induced blood pressure reduction may be attributed to its beneficial effects on the autonomic neurological function. Impaired baroreflex sensitivity has been increasingly postulated to be one of the major causative factors of essential hypertension. Practice of yogic postures has been shown to restore baroreflex sensitivity.<sup>[12,15,31]</sup> Level of adverse neurohormonal activity can be quantitated by the measurement of specific markers in serum and urine. It has been described that regular practitioners of yoga asanas showed a significant reduction in the markers of intrinsic neurohormonal activity such as urinary excretion of adrenaline, noradrenaline, dopamine, aldosterone, as well as serum testosterone and luteinizing hormone levels.<sup>[17,32]</sup>

It is known that the mental state can influence the hypothalamic-pituitary-adrenal (HPA) axis and alter the cortisol levels. Cortisol, discharged from the adrenal cortex, is a biochemical index of HPA axis activation and is a significant marker associated with psychological stress.<sup>[33,34]</sup>

Yoga promotes a reduction in sympathetic activation, enhancement of cardiovagal function, and a shift in autonomic nervous system from primarily sympathetic to parasympathetic,<sup>[17,32]</sup> increased vagal tone, decreased workload on heart, decreased cardiac output, and decreased SBP.<sup>[2]</sup> Asanas such as paschimottanasana have been proved to increase baroreceptor sensitivity.<sup>[33]</sup> Yogic practice, through the restoration of

baroreceptor sensitivity, resulted in significant reduction in the blood pressure of patients who participated in yoga exercise.<sup>[2,26,32]</sup> Both mental relaxation and slow breathing results in a fall in systolic blood pressure, diastolic blood pressure, heart rate, and respiratory rate.<sup>[35]</sup>

**Diastolic blood pressure (mmHg):** In accordance to our finding, van Montfrans et al.,<sup>[36]</sup> Steptoe,<sup>[27]</sup> Patel and Marmot,<sup>[28]</sup> Wenneberg et al.,<sup>[29]</sup> and Joseph et al.<sup>[25]</sup> showed significant findings. Yoga alters hypothalamic discharge leading to decrease in sympathetic tone, peripheral resistance, and, hence, DBP.<sup>[2,13]</sup> Meditation practice results in decreased sympathetic activation, decreased hypothalamic-pituitary-adrenocortical activation, enhanced neurophysiological function, and enhanced serotonin metabolism. Such global physiological changes contribute to the reduction in BP seen across all sex and risk factor subgroups.<sup>[22]</sup>

**ART and VRT:** In accordance to our finding, studies done by Malathi and Parulkar,<sup>[10]</sup> Madanmohan et al.,<sup>[37]</sup> Bhavanani et al.,<sup>[38]</sup> Noorjahan et al.,<sup>[39]</sup> and Jore et al.<sup>[40]</sup> showed statistically significant reduction in ART values after yoga practice. Mean VRT values in normal subjects reduced from 196.17 ± 5.28 milliseconds on the 1st day to 185.25 ± 4.72 **milliseconds after 3 months**. Mean VRT values in hypertensive subjects reduced from 192.25 ± 8.33 milliseconds on the 1st day to 176.35 ± 6.76 milliseconds after 3 months. Results are statistically significant.

In accordance to our finding, studies done by Malathi and Parulkar,<sup>[10]</sup> Madanmohan et al.,<sup>[37]</sup> Bhavanani et al.,<sup>[38]</sup> Noorjahan et al.,<sup>[39]</sup> and Jore et al.<sup>[40]</sup> showed statistically significant reduction in VRT values after yoga practice.

Regular practice of meditation is linked with increased thickness in a subset of cortical regions associated with somatosensory, auditory, visual, and interoceptive processing.<sup>[41,42]</sup> Activation of right hemisphere is seen during the actual act of meditation.<sup>[41,42]</sup>

It is especially fascinating to find that a lot of changes linked with the active meditation state were noted in the frontal lobe. This area is involved with the focus of attention.<sup>[17]</sup> Regular meditation practice may bring down age-related thinning of the frontal cortex. Increased cortical thickness could be owing to greater arborization per neuron, increased glial volume, or raised regional vasculature.<sup>[21,40,41]</sup> The right anterior insula is associated with bodily attention and raised visceral awareness. Increased thickness in this region is associated with increased capacity for awareness.<sup>[43]</sup>

The right hemisphere is essential for sustaining attention, which is a center for insight meditation. It has been hypothesized that by becoming increasingly more aware of sensory stimuli during formal practice, the meditation practitioner is progressively able to use this self-awareness to more successfully navigate through potentially stressful encounters that emerge throughout the day.<sup>[44,45]</sup> Long-term meditation practice is linked with modified resting electroencephalogram patterns, indicative of long-lasting alterations in brain activity.<sup>[42]</sup> Meditation reflects cognitive brain functions such as sequential information processing, stimulus discrimination, and short-term memory.<sup>[46]</sup>

Yoga reduces physiological arousal, simultaneously improving performance on tasks requiring attention.<sup>[47-50]</sup>

## CONCLUSION

It was found that changes in the physiological parameters were significant in hypertensive subjects when compared with normal individuals. Yoga is a helpful intervention in hypertensive subjects. Yoga and pranayam are more beneficial to hypertensive subjects than normal individuals.

In normal subjects, yoga is helpful for keeping the parameters at the baseline, which plays a role in prevention of the disease.

Pranayam and meditation improve (a) sensory information processing ability; (b) central integration of learning and memory, and (c) motor function and coordination, visual scanning, mental flexibility, sustained attention, psychomotor speed, and speed of information processing.<sup>[49,50]</sup> These skills improves sustained attention. This probably improves ART and VRT. This is of applied value in conditions needing faster reactivity such as sports, machine operation, race driving, and specialized surgery. It may also be of value to train mentally challenged children and older sports persons who have prolonged RT. Moreover, yoga and pranayam reduce weight, pulse rate, and blood pressure, which are the risk factors for noncommunicable diseases.

## REFERENCES

1. Sparrenberger F, Fuchs SC, Moreira LB, Fuchs FD. Stressful life events and current psychological distress are associated with self-reported hypertension but not with true hypertension: results from a cross-sectional population-based study. *BMC Public Health*. 2008;8:357.
2. Dusek JA, Hibberd PL, Buczynski B, Chang BH, Dusek KC, Johnston JM, et al. Stress management versus lifestyle modification on systolic hypertension and medication elimination: a randomized trial. *J Altern Complement Med*. 2008;14(2):129-38.
3. Bali LR. Long-term effect of relaxation on blood pressure and anxiety levels of essential hypertensive males: a controlled study. *Psychosom Med*. 1979;41(8):637-46.
4. Khadka R, Paudel BH, Sharma VP, Kumar S, Bhattacharya N. Effect of yoga on cardiovascular autonomic reactivity in essential hypertensive patients. *Health Renaiss*. 2010;8(2):102-9.
5. Hokanson JE, Austin MA. Plasma triglyceride level is a risk factor for cardiovascular disease independent of high-density lipoprotein cholesterol level: a meta-analysis of population-based prospective studies. *J Cardiovasc Risk*. 1996;3(2):213-9.
6. Relation between overweight, diabetes, stress and hypertension: a case-control study in Yarumal-Antioquia, Colombia. *Eur J Epidemiol*. 2001;17(3):275-80.
7. Palaniappan L, Simons LA, Simons J, Friedlander Y, McCallum J. Comparison of usefulness of systolic, diastolic, and mean blood pressure and pulse pressure as predictors of cardiovascular death in patients ≥60 years of age (The Dubbo Study). *Am J Cardiol*. 2002;90(12):1398-1401.
8. Sengupta P. Health impacts of yoga and pranayama: A state-of-the-Art Review. *Int J Prev Med*. 2012;3(7):444-58.



9. Andrews G, Macmohan SW, Austin A, Byrne DG. Hypertension: comparison of drug and non-drug treatments. *Br Med J (Clin Res Ed)*. 1982;284(6328):1523-6.
10. Malathi A, Parulkar VG. Effect of yogasanas on the visual and auditory reaction time. *Indian J Physiol Pharmacol*. 1989;33(2):110-2.
11. Tundwala V, Gupta RP, Kumar S, Singh VB, Sandeep BR, Dayal P, Prakash P. A Study on effect of yoga and various asanas on obesity, hypertension and dyslipidemia. *Int J Basic Appl Med Sci*. 2012;2(1):93-8.
12. Jaysinghe SR. Yoga in cardiac health (a review). *Eur J Cardiovasc Prev Rehabil*. 2004;11(5):369-75.
13. Bharshankar JR, Bharshankar RN, Deshpande VN, Kaore SB, Gosavi GB. Effect of yoga on cardiovascular system in subjects above 40 years. *Indian J Physiol Pharmacol*. 2003;47(2):202-6.
14. Vijayalakshmi P, Madanmohan, Bhavanani AB, Patil A, Babu K. Modulation of stress induced by isometric handgrip test in hypertensive patients following yogic relaxation training. *Indian J Physiol Pharmacol*. 2004;48(1):59-64.
15. Bhavanani AB, Madanmohan, Sanjay Z. Immediate effect of sukha pranayama on cardiovascular variables in patients of hypertension. *Int J Yoga Therap*. 2011;(21):73-6.
16. Gordon L, Morrison EY, McGrowder DA, Young RE, Garwood D, Zamora EM, et al. Changes in clinical and metabolic parameters after exercise therapy in patients with type 2 diabetes. *Arch Med Sci*. 2008;4(4):427-37.
17. Bussing A, Michalsen A, Khalsa SB, Telles S, Sherman KJ. Effects of yoga on mental and physical health: A short summary of reviews. *Evid Based Comp Alternat ed*. 2012;2012:165410.
18. Raub JA. Psychophysiological effects of Hatha Yoga on musculoskeletal and cardiopulmonary function: a literature review. *J Altern Complement Med*. 2002;8(6):797-812.
19. Doijad V, Surdi A. Effect of short term yoga practices on cardiorespiratory fitness parameters. *Int J Basic Med Sci*. 2012;3(5):38-41.
20. Abel AN, Lloyd LK, Williams JS, Miller BK. Physiological characteristics of long-term Bikram yoga practitioners. *J Exerc Physiol Online*. 2012;15(5):32-9.
21. Sundar S, Agrawal SK, Singh VP, Bhattacharya SK, Udupa KN, Vaish SK. Role of yoga in management of essential hypertension. *Acta Cardiol*. 1984;39(3):203-8.
22. Murugesan R, Govindarajulu N, Bera TK. Effect of selected yogic practices on the management of hypertension. *Indian J Physiol Pharmacol*. 2000;44(2):207-10.
23. Dickinson H, Beyer F, Ford G, Nicolson DJ, Campbell F, Cook J, et al. Relaxation therapies for the management of primary hypertension in adults: a Cochrane review. *J Hum Hypertens* 2008;22(12):809-20.
24. Schmidt T, Wijga A, Von Zur Muhlen A, Brabant G, Wagner TO. Changes in cardiovascular risk factors and hormones during a comprehensive residential three month kriya yoga training and vegetarian nutrition. *Acta Physiol Scand Suppl*. 1997;640:158-62.
25. Joseph CN, Porta C, Casucci G, Casiraghi N, Maffei M, Rossi M, et al. Slow breathing improves arterial baroreflex sensitivity and decreases blood pressure in essential hypertension. *Hypertension*. 2005;46:714-8.
26. Alexander CN, Schneider RH, Staggers F, Sheppard W, Clayborne BM, Rainforth M, et al. Trial of stress reduction for hypertension in older African Americans. II. Sex and risk subgroup analysis. *Hypertension*. 1996;28(2):228-37.
27. Steptoe A. Psychological methods in treatment of hypertension: a review. *Br Heart J*. 1977;39(6):587-93.
28. Patel C, Marmot M. Can general practitioners use training in relaxation and management of stress to reduce mild hypertension? *Br Med J* 1988;296(6614):21-4.
29. Wenneberg SR, Schneider RH, Walton KG, Maclean CR, Levitsky DK, Salerno JW, et al. A controlled study of the effects of the Transcendental Meditation program on cardiovascular reactivity and ambulatory blood pressure. *Int J Neurosci*. 1997;89(1-2):15-28.
30. Granath J, Ingvarsson S, von Thiele U, Lundberg U. Stress management: a randomized study of cognitive behavioural therapy and yoga. *Cogn Behav Ther*. 2006;35(1):3-10.
31. Gordon L, Morrison EY, McGrowder D, Penas YF, Zamoraz EM, Garwood D, et al. Effect of yoga and traditional physical exercise on hormones and percentage insulin binding receptor in patients with type 2 diabetes. *Am J Biochem Biotechnol*. 2008;4(1):35-42.
32. Innes KE, Selfe TK, Alexander GK. The Effects of a Gentle Yoga Program on Sleep, Mood, and Blood Pressure in Older Women with Restless Legs Syndrome (RLS): A Preliminary Randomized Controlled Trial. *Evid Based Complement Alternat Med*. 2012;2012:294058.
33. Yoshihara K, Hiramoto T, Sudo N, Kubo C. Profile of mood states and stress-related biochemical indices in long-term yoga practitioners. *Biopsychosoc Med*. 2011;5(1):5-6.
34. Shantakumari N, Sequeira S, Eldeeb R. Effect of a yoga intervention on hypertensive diabetic patients. *J Adv Int Med*. 2012;1(2):60-3.
35. Satyapriya M, Nagendra H R, Nagarathna R, Padmalatha V. Effect of integrated yoga on stress and heart rate variability in pregnant women. *Int J Gynaecol Obstet*. 2009;104(3):218-22.
36. van Montfrans GA, Karemaker JM, Wieling W, Dunning AJ. Relaxation therapy and continuous ambulatory blood pressure in mild hypertension: a controlled study. *BMJ*. 1990;300(6736):1368-72.
37. Madanmohan, Udupa K, Bhavanani AB, Shatapathy CC, Sahai A. Modulation of cardiovascular response to exercise by yoga training. *Indian J Physiol Pharmacol*. 2004;48(4):461-5.
38. Bhavanani AB, Ramanathan M, Kt H. Immediate effect of mukha bhastrika (a bellows type pranayama) on reaction time in mentally challenged adolescents. *Indian J Physiol Pharmacol*. 2012;56(2):174-80.
39. Begum N, Kumaran SD, Venkatesh G, Kulkarni SB. Study of the utility of short course of yoga to improve reaction time. *J Clin Diagn Res*. 2012;6(7):1241-3.
40. Jore SB, Kamble P, Bhutada TB, Patwardhan MS. Effect of pranayama training on audio-visual reaction time. *Int J Healthc Biomed Res*. 2013;2(1):35-7.
41. Lazar SW, Kerr CE, Wasserman RH, Gray JR, Greve DN, Treadway MT, et al. Meditation experience is associated with increased cortical thickness. *Neuroreport*. 2005;16(17):1893-7.
42. Cohen DL, Wintering N, Tolles V, Townsend RR, Farrar JT, Galantino ML, et al. Cerebral blood flow effects of yoga training: preliminary evaluation of 4 cases. *J Altern Complement Med*. 2009;15(1):9-14.
43. Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting interoceptive awareness. *Nat Neurosci*. 2004;7(2):189-95.
44. Goldstein J, Kornfield J. *Seeking the Heart of Wisdom: The Path of Inside Meditation*. Boston: Shambhala Publication, 1987.
45. Craig AD. Interoception: the sense of the physiological condition of the body. *Curr Opin Neurobiol*. 2003;13(4):500-5.
46. Ospina MB, Bond K, Karkhaneh M, Tjosvold L, Vandermeer B, Liang Y, et al. Meditation practices for health: state of the research. *Evid Rep Technol Assess (Full Rep)*. 2007;(155):1-263.

47. Bijlani RL. Influence of yoga on brain and behavior: facts and speculations. *Indian J Physiol Pharmacol.* 2004;48(1):1-5.
48. Madanmohan, Udupa K, Bhavanani AB, Vijayalakshmi P, Surendiran A. Effect of slow and fast pranayams on reaction time and cardio-respiratory variables. *Indian J Physiol Pharmacol.* 2005;49(3): 313-8.
49. Sharma V, Kukreja A, Senthil Kumar S, Kanojia S, Gupta S, Das S, et al. Comparative study of yoga and physical exercises on psychological parameters, hand grip strength and reaction time during examination stress in female medical students. *Int J Publ Health Res Dev.* 2012;2(1):362.
50. Dash M, Telles S. Yoga training and motor speed based on a finger tapping task. *Indian J Physiol Pharmacol.* 1999;43(4):458-62.

**How to cite this article:** Sonwane TD, Mishra NV. Effects of yoga and pranayam on human reaction time and certain physiological parameters in normal and hypertensive subjects. *Natl J Physiol Pharm Pharmacol* 2016;6:323-328

**Source of Support:** Nil, **Conflict of Interest:** None declared.